



Diabetes Research Open Access: Sodium Glucose Cotransporter2 (SGLT2) Inhibitors

Kuşkonmaz SM^{1*}

¹Ankara Training and Research Hospital, Department of Endocrinology and Metabolism, Ulucanlar Street No 89, 06230, Turkey

Corresponding Author: **Şerife Mehlika Kuşkonmaz**

Address: Ankara Training and Research Hospital, Department of Endocrinology and Metabolism, Ulucanlar Street No 89, 06230, Turkey; Email: mehlikaisildak@gmail.com

Received date: 28 January 2020; **Accepted date:** 01 February 2020; **Published date:** 08 February 2020

Citation: Kuşkonmaz SM. Diabetes Research Open Access: Sodium Glucose Cotransporter2 (SGLT2) Inhibitors. Diab Res Open Access. 2020 Feb 08;2(S1):14.

Copyright © 2020 Kuşkonmaz SM. This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords

Diabetes Research; Sodium Glucose Cotransporter2 (SGLT2) Inhibitors; Special Issue

Sodium glucose cotransporter 2 (SGLT2) inhibitors (SGLT2i) are a group of glycosuric drugs approved in the management of type 2 diabetes mellitus. They act on the sodium glucose cotransporter and inhibit renal glucose reabsorption. Canagliflozin dapagliflozin and empagliflozin are members of the SGLT2i group. SGLT2 is supposed to be unique to the kidney. Recent studies showed the benefits of these agents beyond and independent from glucose lowering. New guidelines emphasize these pleiotropic effects such as cardioprotective and renoprotective effects of SGLT2i and suggest them as first line oral antidiabetics in patients with coronary heart disease.

On the other hand, adverse events related to SGLT2i are being reported in the literature. SGLT2i are found to be associated with euglycemic ketoacidosis, fractures and acute kidney injury. Food and Drug Administration (FDA) recently added a warning about the risk of serious genital infections with SGLT2i to the previous warning about the risk of leg and foot amputations specifically with

canagliflozin.

In this issue of Diabetes Research, we wanted to evaluate these novel agents with all aspects and aimed to summarize the beneficial effects of and possible adverse events related to SGLT2i. We hope the clinicians who deal with diabetes will benefit much from the concise reading of this special issue of Diabetes Research.

