



## Latest Standard Management for Heart Failure with Guideline-Directed Medical Therapy (GDMT)

Hiroshi BANDO<sup>1,2\*</sup>

<sup>1</sup>Tokushima University / Medical Research, Tokushima, Japan

<sup>2</sup>Integrative Medicine Japan (IMJ), Shikoku Island division, Tokushima, Japan

Corresponding Author: **Hiroshi BANDO, MD, PhD, FACP** [ORCID ID](#)

**Address:** Tokushima University /Medical Research, Nakashowa 1-61, Tokushima 770-0943, Japan; Tel: +81-90-3187-2485; Email: [pianomed@bronze.ocn.ne.jp](mailto:pianomed@bronze.ocn.ne.jp)

**Received date:** 15 July 2022; **Accepted date:** 16 August 2022; **Published date:** 20 August 2022

**Citation:** Bando H. Latest Standard Management for Heart Failure with Guideline-Directed Medical Therapy (GDMT). J Health Care and Research. 2022 Aug 20;3(2):35-40.

**Copyright** © 2022 Bando H. This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium provided the original work is properly cited.

### Abstract

Sodium–glucose cotransporter 2 inhibitor (SGLT2i) has been attracting attention for novel agent for patients with diabetes and also heart failure (HF), in which the left ventricular ejection fraction (LVEF) has decreased. For standard cardiovascular treatment, 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure is recently presented. Some important perspectives were found, which are HF with mildly reduced EF (HFmrEF), HF with improved EF (HFimpEF) and HF with preserved EF (HFpEF). For patients with HFmrEF, SGLT2i can contribute reducing HF hospitalizations and cardiovascular death. From now, the guideline-directed medical therapy (GDMT) will contribute the standard and beneficial therapy.

### Keywords

Sodium–Glucose Cotransporter 2 Inhibitor, Left Ventricular Ejection Fraction, Heart Failure with Mildly Reduced EF, Guideline-Directed Medical Therapy, 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure

### Abbreviations

ACEi: Angiotensin-Converting Enzyme Inhibitor; ARB: Angiotensin Receptor Blocker; ARNi: Angiotensin Receptor–Neprilysin Inhibitor; COR: Class of Recommendation; GDMT: Guideline-Directed Medical Therapy; HF: Heart Failure; HFmrEF: Heart Failure with Mildly Reduced Ejection Fraction; HFimpEF: Heart Failure with Improved Ejection Fraction; HFpEF: Heart Failure with Preserved Ejection Fraction; LVEF: Left Ventricular Ejection Fraction; MRA: Mineralocorticoid Receptor Antagonist; SGLT2i: Sodium-Glucose Cotransporter 2 Inhibitor; CANVAS: Choice of ANesthesia for EndoVascular Treatment of Acute Ischemic Stroke; DECLARE-TIMI 58 trial: Dapagliflozin Effect on Cardiovascular Events – Thrombolysis in Myocardial Infarction 58; EMPEROR-Reduced: Empagliflozin in Patients with Chronic Heart Failure and a Reduced Ejection Fraction; DAPA-HF: The Dapagliflozin and Prevention of Adverse-Outcomes in Heart Failure (DAPA-HF) Trial; EMPA-REG OUTCOME: (Empagliflozin) Cardiovascular Outcome Event Trial in Type 2 Diabetes Mellitus Patients (EMPA-REG OUTCOME) study

Sodium-glucose cotransporter 2 inhibitor (SGLT2i) was originally developed as a drug for blood glucose control [1,2]. Data from clinical trials have shown that it is also useful for controlling heart failure events in diabetic patients [3,4]. For these trials, three representative studies were observed including EMPA-REG OUTCOME, CANVAS and DECLARE-TIMI 58. From clinical trials in patients with heart failure, major adverse cardiovascular events (MACE) and heart failure events were suppressed by SGLT2i regardless of the presence of diabetes. For the related studies, EMPEROR-Reduced and DAPA-HF were included. SGLT2i is now attracting attention as a new therapeutic agent for patients with heart failure (HF), in which the left ventricular ejection fraction (LVEF) has decreased.

From a historical point of view, the standard therapy for heart failure has been changed for decades. In 2013, ACCF/AHA Guideline for the Management of Heart Failure was observed and used at that time [5]. Successively, in 2017, ACC/AHA/HFSA focused update of the 2013 edition was found [6]. The latest topic would be the presentation of the latest announcement of revision, that is the "2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure". Its characteristic points mean the intention for providing patient-centered recommendations of clinicians in the actual medical practice [7]. They include prevention, diagnosis and management for many patients with heart failure.

For the preparation of the standard guideline, lots of literature were searched from the period May-Dec 2020 [7]. The fields included reviews, various studies and other reports of English language which were from EMBASE, PubMed (MEDLINE), the Agency for Healthcare Research and Quality, Cochrane Collaboration and other databases. Furthermore, other relevant clinical and research investigations during Jan-Sept 2021 have been considered for analysis. The current standard guideline has been harmonized and collaborated with other main medical associations, including American College of Cardiology (ACC) and American Heart Association (AHA) for the period until Dec 2021.

For long years, heart failure has been the most prevalent cause of mortality and morbidity across the world. Current 2022 edition of the guideline for heart failure can provide adequate recommendations which are contemporary evidence-based aspects for actual treatment of HF patients [7]. These recommendations in the guideline can show some evidence-based approach method for applicable managing cases with heart failure. It includes the direction for improving the quality of care and cure along with each interest of the case. Former recommendations in the previous edition were updated and changed based on new evidence. Furthermore, recent recommendations have been announced that were strengthened by published evidence. With reported economic analysis, value statements were also included for several methods of treatment.

This guideline was proposed by the combination of ACC/AHA/HFSA, and its executive summary includes several crucial points [8]. There are four class of recommendation (COR) for the patients, where COR A: At-risk for heart failure (HF), B: pre-HF, C: symptomatic HF and D: advanced HF. COR A means the patients with hypertension, diabetes and others with possible arteriosclerosis without current or previous symptoms/signs of HF. The progress is found from COR A, B, C and D in this order.

In actual clinical practice, the following perspectives are beneficial from medical, cardiovascular points of view. They are i) there are Guideline-directed medical therapy (GDMT) for heart failure (HF) with reduced ejection fraction (HFrEF), ii) it has 4 categories of medications including SGLT2i, iii) The four groups reveal a) RAS inhibition with ARNi, ACEi, ARB, b) beta-blockers, c) MRAs, d) new group that is SGLT2i. iv) SGLT2i shows a class of recommendation 2a for HFmrEF, and weaker recommendation 2b are revealed for beta-blocker, ACEi, ARB, MRA and ARNi, v) New recommendations for HFpEF are for SGLT2i (2a), MRAs and ARNi (2b), vi) Improved LVEF has been used to refer to HF patient with previous HFrEF who shows now > 40% of LVEF, and such case has to continue HFrEF treatment, vii) When diagnosing HF with > 40% of LVEF, supporting the evidence of elevated filling pressure is important, viii) for

obtaining the evidence, invasive or noninvasive biomarkers can be beneficial such as hemodynamic exam, diastolic function with imaging or natriuretic peptide measurement [8].

In the current AHA/ACC/HFSA guideline for the management of heart failure, the most impressive point would be the following judgment. Treatment of HFmrEF (LVEF 41-49%) and HFpEF (LVEF > 50%) showed common evidence levels, which are diuretics as needed (1), SGLT2i (2a)[9] and ARNi[10] /MRA[11] / ARB (2b)[12,13]. The former only showed ACEi (2b)[14] and evidence-based beta blockers for HFrEF (2b)[15]. SGLT2i has been highly evaluated for HF, and it shows 1 for COR and A for Level of Evidence (LOE). In the case of symptomatic chronic HFrEF, SGLT2i agent has been recommended to decrease the hospitalization for HF and cardiovascular mortality, that is with or without T2D presence from DAPA-HF and EMPEROR-Reduced [9,16].

In the latest guideline, some important perspectives were found, which are a) Heart Failure with mildly reduced EF (HFmrEF), b) HF with improved EF (HFimpEF) and c) HF with preserved EF (HFpEF). For patients with HFmrEF, SGLT2i can contribute to reducing HF hospitalizations and cardiovascular death [17]. In the case of 41-49% of LVEF, HF hospitalizations/CV death can be decreased by ACEi, ARB, MRA, ARNi, beta-blocker [18,19]. Cases with HFmrEF would have repeated LVEF evaluation for clinical progress. For patients with HFimpEF after treatment, GDMT would be followed up for preventing LV dysfunction and HF relapse, even if the case keeps the condition without any symptoms [19].

The condition of HFpEF (LVEF 50%) has been rather prevalent, that seems to be at most 50% of all HF patients and shows significant association of mortality and morbidity [20]. HFpEF reveals a heterogenous situation associated with comorbidities such as diabetes, hypertension, obesity, CKD and others [21]. The definition of HFpEF has been variable in lots of clinical trials [22]. For recent trials, beneficial HFpEF results were not found for HF

hospitalizations [23,24].

In summary, "2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure" has been recently presented, in which SGLT2i shows novel therapeutic agents for HF. HF patients can be categorized as HFmrEF, HFimpEF and HFpEF, where SGLT2i may contribute to decreasing HF hospitalizations and cardiovascular death. This article hopefully becomes a useful reference for cardiovascular and diabetic practice and research.

### Conflict of Interest

The author has read and approved the final version of the manuscript. The author has no conflicts of interest to declare.

### References

- [1] Bhattarai M, Salih M, Regmi M, Al-Akchar M, Deshpande R, Niaz Z, Kulkarni A, Siddique M, Hegde S. Association of Sodium-Glucose Cotransporter 2 Inhibitors With Cardiovascular Outcomes in Patients With Type 2 Diabetes and Other Risk Factors for Cardiovascular Disease: A Meta-analysis. *JAMA Netw Open.* 2022 Jan 4;5(1):e2142078. [PMID: 34985519]
- [2] Salah HM, Al'Aref SJ, Khan MS, Al-Hawwas M, Vallurupalli S, Mehta JL, Mounsey JP, Greene SJ, McGuire DK, Lopes RD, Fudim M. Efficacy and safety of sodium-glucose cotransporter 2 inhibitors initiation in patients with acute heart failure, with and without type 2 diabetes: a systematic review and meta-analysis. *Cardiovasc Diabetol.* 2022 Feb 5;21(1):20. [PMID: 35123480]
- [3] Zou X, Shi Q, Vandvik PO, Guyatt G, Lang CC, Parpia S, Wang S, Agarwal A, Zhou Y, Zhu Y, Tian H, Zhu Z, Li S. Sodium-Glucose Cotransporter-2 Inhibitors in Patients With Heart Failure: A Systematic Review and Meta-analysis. *Ann Intern Med.* 2022 Jun;175(6):851-61. [PMID: 35404670]
- [4] Tomasoni D, Fonarow GC, Adamo M, Anker SD, Butler J, Coats AJS, Filippatos G, Greene SJ, McDonagh TA, Ponikowski P, Rosano G, Seferovic P, Vaduganathan M, Voors AA, Metra M. Sodium-glucose co-transporter 2 inhibitors as an early, first-line therapy in patients with heart failure and reduced

- ejection fraction. *Eur J Heart Fail.* 2022 Mar;24(3):431-41. [PMID: 34894038]
- [5] Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL; American College of Cardiology Foundation; American Heart Association Task Force on Practice Guidelines. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2013 Oct 15;62(16):e147-39. [PMID: 23747642]
- [6] Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos GS, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation.* 2017 Aug 8;136(6):e137-61. [PMID: 28455343]
- [7] Writing Committee Members; ACC/AHA Joint Committee Members. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure. *J Card Fail.* 2022 May;28(5):e1-67. [PMID: 35378257]
- [8] Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, Fang JC, Fedson SE, Fonarow GC, Hayek SS, Hernandez AF, Khazanie P, Kittleson MM, Lee CS, Link MS, Milano CA, Nwacheta LC, Sandhu AT, Stevenson LW, Vardeny O, Vest AR, Yancy CW; WRITING COMMITTEE MEMBERS. 2022 American College of Cardiology/American Heart Association/Heart Failure Society of America Guideline for the Management of Heart Failure: Executive Summary. *J Card Fail.* 2022 May;28(5):810-30. [PMID: 35378259]
- [9] Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, Januzzi J, Verma S, Tsutsui H, Brueckmann M, Jamal W, Kimura K, Schnee J, Zeller C, Cotton D, Bocchi E, Böhm M, Choi DJ, Chopra V, Chuquiure E, Giannetti N, Janssens S, Zhang J, Gonzalez Juanatey JR, Kaul S, Brunner-La Rocca HP, Merkely B, Nicholls SJ, Perrone S, Pina I, Ponikowski P, Sattar N, Senni M, Seronde MF, Spinar J, Squire I, Taddei S, Wanner C, Zannad F; EMPEROR-Reduced Trial Investigators. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med.* 2020 Oct 8;383(15):1413-24. [PMID: 32865377]
- [10] Wang Y, Zhou R, Lu C, Chen Q, Xu T, Li D. Effects of the Angiotensin-Receptor Neprilysin Inhibitor on Cardiac Reverse Remodeling: Meta-Analysis. *J Am Heart Assoc.* 2019 Jul 2;8(13):e012272. [PMID: 31240976]
- [11] Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ, Pitt B; EMPHASIS-HF Study Group. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med.* 2011 Jan 6;364(1):11-21. [PMID: 21073363]
- [12] Desai AS, Solomon SD, Shah AM, Claggett BL, Fang JC, Izzo J, McCague K, Abbas CA, Rocha R, Mitchell GF; EVALUATE-HF Investigators. Effect of Sacubitril-Valsartan vs Enalapril on Aortic Stiffness in Patients with Heart Failure and Reduced Ejection Fraction: A Randomized Clinical Trial. *JAMA.* 2019 Sep 17;322(11):1077-84. [PMID: 31475296]
- [13] Konstam MA, Neaton JD, Dickstein K, Drexler H, Komajda M, Martinez FA, Riegger GA, Malbecq W, Smith RD, Gupta S, Poole-Wilson PA; HEAAL Investigators. Effects of high-dose versus low-dose losartan on clinical outcomes in patients with heart failure (HEAAL study): a randomised, double-blind trial. *Lancet.* 2009 Nov 28;374(9704):1840-48. Erratum in: *Lancet.* 2009 Dec 5;374(9705):1888. [PMID: 19922995]
- [14] Telmisartan Randomised Assessment Study in ACE Intolerant subjects with cardiovascular Disease (TRANSCEND) Investigators, Yusuf S, Teo K, Anderson C, Pogue J, Dyal L, Copland I, Schumacher H, Dagenais G, Sleight P. Effects of the angiotensin-receptor blocker telmisartan on cardiovascular events in high-risk patients intolerant to angiotensin-converting enzyme inhibitors: a randomised controlled

trial. *Lancet*. 2008 Sep 27;372(9644):1174-83. Erratum in: *Lancet*. 2008 Oct 18;372(9647):1384. [PMID: 18757085]

[15] Packer M, Fowler MB, Roecker EB, Coats AJ, Katus HA, Krum H, Mohacsi P, Rouleau JL, Tendera M, Staiger C, Holcslaw TL, Amann-Zalan I, DeMets DL; Carvedilol Prospective Randomized Cumulative Survival (COPERNICUS) Study Group. Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the carvedilol prospective randomized cumulative survival (COPERNICUS) study. *Circulation*. 2002 Oct 22;106(17):2194-99. [PMID: 12390947]

[16] McMurray JJV, Solomon SD, Inzucchi SE, Køber L, Kosiborod MN, Martinez FA, Ponikowski P, Sabatine MS, Anand IS, Bělohávek J, Böhm M, Chiang CE, Chopra VK, de Boer RA, Desai AS, Diez M, Drozd J, Dukát A, Ge J, Howlett JG, Katova T, Kitakaze M, Ljungman CEA, Merkely B, Nicolau JC, O'Meara E, Petrie MC, Vinh PN, Schou M, Tereshchenko S, Verma S, Held C, DeMets DL, Docherty KF, Jhund PS, Bengtsson O, Sjöstrand M, Langkilde AM; DAPA-HF Trial Committees and Investigators. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med*. 2019 Nov 21;381(21):1995-2008. [PMID: 31535829]

[17] Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, Brunner-La Rocca HP, Choi DJ, Chopra V, Chuquiure-Valenzuela E, Giannetti N, Gomez-Mesa JE, Janssens S, Januzzi JL, Gonzalez-Juanatey JR, Merkely B, Nicholls SJ, Perrone SV, Piña IL, Ponikowski P, Senni M, Sim D, Spinar J, Squire I, Taddei S, Tsutsui H, Verma S, Vinereanu D, Zhang J, Carson P, Lam CSP, Marx N, Zeller C, Sattar N, Jamal W, Schnaidt S, Schnee JM, Brueckmann M, Pocock SJ, Zannad F, Packer M; EMPEROR-Preserved Trial Investigators. Empagliflozin in Heart Failure with a Preserved Ejection Fraction. *N Engl J Med*. 2021 Oct 14;385(16):1451-61. [PMID: 34449189]

[18] Solomon SD, McMurray JJV, Anand IS, Ge J, Lam CSP, Maggioni AP, Martinez F, Packer M, Pfeffer MA, Pieske B, Redfield MM, Rouleau JL, van Veldhuisen DJ, Zannad F, Zile MR, Desai AS, Claggett B, Jhund PS, Boytsov SA, Comin-Colet J, Cleland J, Düngen HD, Goncalvesova E, Katova T, Kerr Saraiva JF, Lelonek M, Merkely B, Senni M, Shah SJ, Zhou J,

Rizkala AR, Gong J, Shi VC, Lefkowitz MP; PARAGON-HF Investigators and Committees. Angiotensin-Neprilysin Inhibition in Heart Failure with Preserved Ejection Fraction. *N Engl J Med*. 2019 Oct 24;381(17):1609-20. [PMID: 31475794]

[19] Halliday BP, Wassall R, Lota AS, Khaliq Z, Gregson J, Newsome S, Jackson R, Rahneva T, Wage R, Smith G, Venneri L, Tayal U, Auger D, Midwinter W, Whiffin N, Rajani R, Dzungu JN, Pantazis A, Cook SA, Ware JS, Baksi AJ, Pennell DJ, Rosen SD, Cowie MR, Cleland JGF, Prasad SK. Withdrawal of pharmacological treatment for heart failure in patients with recovered dilated cardiomyopathy (TRED-HF): an open-label, pilot, randomised trial. *Lancet*. 2019 Jan 5;393(10166):61-73. [PMID: 30429050]

[20] Cincin A, Abul Y, Ozben B, Tanrikulu A, Topaloglu N, Ozgul G, Karakurt S, Oktay A. Pleural fluid aminoterminal brain natriuretic peptide in patients with pleural effusions. *Respir Care*. 2013 Feb;58(2):313-19. [PMID: 22710710]

[21] Mentz RJ, Kelly JP, von Lueder TG, Voors AA, Lam CS, Cowie MR, Kjeldsen K, Jankowska EA, Atar D, Butler J, Fiuzat M, Zannad F, Pitt B, O'Connor CM. Noncardiac comorbidities in heart failure with reduced versus preserved ejection fraction. *J Am Coll Cardiol*. 2014 Dec 2;64(21):2281-93. [PMID: 25456761]

[22] Ho JE, Zern EK, Wooster L, Bailey CS, Cunningham T, Eisman AS, Hardin KM, Zampierollo GA, Jarolim P, Pappagianopoulos PP, Malhotra R, Nayor M, Lewis GD. Differential Clinical Profiles, Exercise Responses, and Outcomes Associated With Existing HFpEF Definitions. *Circulation*. 2019 Jul 30;140(5):353-65. [PMID: 31132875]

[23] Solomon SD, Vaduganathan M, L Claggett B, Packer M, Zile M, Swedberg K, Rouleau J, A Pfeffer M, Desai A, Lund LH, Kober L, Anand I, Sweitzer N, Linssen G, Merkely B, Luis Arango J, Vinereanu D, Chen CH, Senni M, Sibulo A, Boytsov S, Shi V, Rizkala A, Lefkowitz M, McMurray JJV. Sacubitril/Valsartan Across the Spectrum of Ejection Fraction in Heart Failure. *Circulation*. 2020 Feb 4;141(5):352-61. [PMID: 31736342]

[24] Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Harty B, Heitner JF, Kenwood CT, Lewis EF, O'Meara E, Probstfield JL, Shaburishvili

Citation: Bando H. Latest Standard Management for Heart Failure with Guideline-Directed Medical Therapy (GDMT). J Health Care and Research. 2022 Aug 20;3(2):35-40.

Commentary

T, Shah SJ, Solomon SD, Sweitzer NK, Yang S, McKinlay SM; TOPCAT Investigators. Spironolactone for heart failure with preserved ejection fraction. N Engl J Med. 2014 Apr 10;370(15):1383-92. [PMID: [24716680](#)]



Keywords: Sodium-Glucose Cotransporter 2 Inhibitor, Left Ventricular Ejection Fraction, Heart Failure with Mildly Reduced EF, Guideline-Directed Medical Therapy, 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure